

Some properties of Kaehler submanifolds with recurrent tensor fields

Irina I. Bodrenko ¹

Abstract

The properties of Kaehler submanifolds with recurrent the second fundamental form in spaces of constant holomorphic sectional curvature are being studied in this article.

Introduction

Let M^{2m+2l} be a Kaehler manifold of complex dimension $m + l$ ($m \geq 1, l \geq 1$) with almost complex structure J and a Riemannian metric $\tilde{g}$, $\tilde{\nabla}$ be the Riemannian connection coordinated with $\tilde{g}$, $\tilde{R}$ be the Riemannian curvature tensor of manifold M^{2m+2l} . Let F^{2m} be a Kaehler submanifold of complex dimension m in M^{2m+2l} with induced Riemannian metric g . The restriction J to F^{2m} defines induced almost complex structure on F^{2m} , which we will denote by the same symbol J . Let ∇ be the Riemannian connection coordinated with g , D be the normal connection, b be the second fundamental form, $R^\perp$ be the tensor of normal curvature of submanifold F^{2m} , $\bar{\nabla} = \nabla \oplus D$ be the connection of van der Waerden — Bortolotti. b is called *parallel* if $\bar{\nabla}b \equiv 0$. A tensor of normal curvature $R^\perp$ is called *parallel* if $\bar{\nabla}R^\perp \equiv 0$.

According to the definition of recurrent tensor field (see [1], note 8), nonzero form $b \neq 0$ is called *recurrent* if there exists 1-form μ on F^{2m} such that $\bar{\nabla}b = \mu \otimes b$.

Theorem 1 . Let F^{2m} be a Kaehler submanifold of complex dimension m in a Kaehler manifold $M^{2m+2l}(c)$ of complex dimension $m+l$ and constant holomorphic sectional curvature c . If F^{2m} has recurrent the second fundamental form b then the tensor of normal curvature $R^\perp \neq 0$ is parallel.

It is known (see [1], note 8, theorem 3), that for a Riemannian manifold M with recurrent tensor of Riemannian curvature $\tilde{R}$ and irreducible narrowed linear group of holonomy, it is necessary that the tensor of Riemannian curvature $\tilde{R}$ be parallel (i.e. $\tilde{\nabla}\tilde{R} \equiv 0$) with the condition $\dim M \geq 3$. A Riemannian manifold M is called *locally symmetric* if $\tilde{\nabla}\tilde{R} \equiv 0$.

Theorem 2 . Let F^{2m} be a Kaehler submanifold of complex dimension m in a Kaehler manifold $M^{2m+2l}(c)$ of complex dimension $m+l$ and constant holomorphic sectional curvature c . If F^{2m} has recurrent the second fundamental form b then F^{2m} is locally symmetric submanifold.

¹©Irina I. Bodrenko, associate professor, Department of Mathematics,
Volgograd State University, University Prospekt 100, Volgograd, 400062, RUSSIA.
E.-mail: bodrenko@mail.ru <http://www.bodrenko.com> <http://www.bodrenko.org>

1 Main notations and formulas.

Let M^{n+p} be $(n + p)$ -dimensional ($n \geq 2, p \geq 2$) smooth Riemannian manifold, $\tilde{g}$ be a Riemannian metric on M^{n+p} , $\tilde{\nabla}$ be the Riemannian connection coordinated with $\tilde{g}$, F^n be n -dimensional smooth submanifold in M^{n+p} , g be the induced Riemannian metric on F^n , ∇ be the Riemannian connection on F^n coordinated with g , TF^n and $T^\perp F^n$ be tangent and normal bundles on F^n , respectively, R and R_1 be the tensors of Riemannian and Ricci curvature of connection ∇ , respectively, b be the second fundamental form F^n , D be the normal connection, $R^\perp$ be the tensor of normal curvature, $\overline{\nabla}$ be the connection of Van der Waerden — Bortolotti.

The formulas of Gauss and Weingarten have, respectively, the following form [2] :

$$\tilde{\nabla}_X Y = \nabla_X Y + b(X, Y), \quad (1.1)$$

$$\tilde{\nabla}_X \xi = -A_\xi X + D_X \xi, \quad (1.2)$$

for any vector fields X, Y , tangent to F^n , and vector field ξ normal to F^n .

The equations of Gauss, Peterson — Codacci and Ricci have, respectively, the following form [2] :

$$\tilde{R}(X, Y, Z, W) = R(X, Y, Z, W) + \tilde{g}(b(X, Z), b(Y, W)) - \tilde{g}(b(X, W), b(Y, Z)), \quad (1.3)$$

$$(\tilde{R}(X, Y)Z)^\perp = (\overline{\nabla}_X b)(Y, Z) - (\overline{\nabla}_Y b)(X, Z), \quad (1.4)$$

$$\tilde{R}(X, Y, \xi, \eta) = R^\perp(X, Y, \xi, \eta) - g([A_\xi, A_\eta]X, Y), \quad (1.5)$$

for any vector fields X, Y, Z, W , tangent to F^n , and vector fields ξ, η , normal to F^n .

For any vector field ξ normal to F^n , we denote as A_ξ the second fundamental tensor relatively to ξ . For A_ξ the relation holds

$$\tilde{g}(b(X, Y), \xi) = \tilde{g}(A_\xi X, Y), \quad (1.6)$$

for any vector fields X, Y , tangent to F^n .

A normal vector field ξ is called *nondegenerate* if $\det A_\xi \neq 0$.

Covariant derivatives $\overline{\nabla}b$, $(\overline{\nabla}A)_\xi$ and $\overline{\nabla}R^\perp$, are defined by the following equalities, respectively ([2]):

$$(\overline{\nabla}_X b)(Y, Z) = D_X(b(Y, Z)) - b(\nabla_X Y, Z) - b(Y, \nabla_X Z), \quad (1.7)$$

$$(\overline{\nabla}_X A)_\xi Y = \nabla_X(A_\xi Y) - A_\xi(\nabla_X Y) - A_{D_X \xi} Y, \quad (1.8)$$

$$(\overline{\nabla}_X R^\perp)(Y, Z)\xi = D_X(R^\perp(Y, Z)\xi) - R^\perp(\nabla_X Y, Z)\xi - R^\perp(Y, \nabla_X Z)\xi - R^\perp(Y, Z)D_X \xi, \quad (1.9)$$

for any vector fields X, Y, Z , tangent to F^n , and vector field ξ normal to F^n .

Let indices, in this article, take the following values: $i, j, k, s, t = 1, \dots, n$, $\alpha, \beta, \gamma = 1, \dots, p$. We will use the Einstein rule.

Let x be an arbitrary point F^n , $T_x F^n$ and $T_x^\perp F^n$ be the tangent and normal spaces F^n at point x , respectively, $U(x)$ be some neighborhood of point x , $(u^1, \dots, u^n)$ be local

coordinates on F^n in $U(x)$, $\{\partial/\partial u^i\}$ be a local basis in TF^n , $\{n_{\alpha|}\}$ be a field of bases of normal vectors in $T^\perp F^n$ in $U(x)$. We may always choose the basis $\{n_{\alpha|}\}$ orthonormalized and assume that $\tilde{g}(n_{\alpha|}, n_{\beta|}) = \delta_{\alpha\beta}$, where $\delta_{\alpha\beta}$ is the Kronecker symbol. We introduce the following designations:

$$\begin{aligned} g_{ij} &= g\left(\frac{\partial}{\partial u^i}, \frac{\partial}{\partial u^j}\right), \quad b_{ij} = b\left(\frac{\partial}{\partial u^i}, \frac{\partial}{\partial u^j}\right) = b_{ij}^\alpha n_{\alpha|}, \quad \Gamma_{ij,k} = g\left(\nabla_{\frac{\partial}{\partial u^i}} \frac{\partial}{\partial u^j}, \frac{\partial}{\partial u^k}\right), \\ \Gamma_{ij}^k &= g^{kt} \Gamma_{ij,t}, \quad \nabla_i b_{jk}^\alpha = \frac{\partial b_{jk}^\alpha}{\partial u^i} - \Gamma_{ij}^t b_{tk}^\alpha - \Gamma_{ik}^t b_{jt}^\alpha, \quad \Gamma_{\alpha\beta|i}^\perp = \tilde{g}\left(n_{\alpha|}, \widetilde{\nabla_{\frac{\partial}{\partial u^i}} n_{\beta|}}\right), \\ \Gamma_{\beta|i}^{\perp\alpha} &= \delta^{\alpha\gamma} \Gamma_{\gamma\beta|i}^\perp, \quad \Gamma_{\alpha\beta|i}^\perp = \delta_{\alpha\gamma} \Gamma_{\beta\gamma|i}^\perp, \quad \Gamma_{\alpha\beta|i}^\perp + \Gamma_{\beta\alpha|i}^\perp = 0, \quad \bar{\nabla}_i b_{jk}^\alpha = \left(\nabla_i b_{jk}^\alpha + \Gamma_{\beta|i}^{\perp\alpha} b_{jk}^\beta\right), \\ \left(\bar{\nabla}_{\frac{\partial}{\partial u^i}} b\right)\left(\frac{\partial}{\partial u^j}, \frac{\partial}{\partial u^k}\right) &= \bar{\nabla}_i b_{jk}^\alpha n_{\alpha|}, \quad b_{\alpha|ik} = \delta_{\alpha\beta} b_{ik}^\beta, \quad a_{\alpha|i}^j = b_{\alpha|ik} g^{kj}, \\ \nabla_i a_{\alpha|j}^k &= \frac{\partial a_{\alpha|j}^k}{\partial u^i} - \Gamma_{ij}^t a_{\alpha|t}^k + \Gamma_{it}^k a_{\alpha|j}^t, \quad a_{\alpha|i}^j \frac{\partial}{\partial u^j} = A_{n_{\alpha|}}\left(\frac{\partial}{\partial u^i}\right), \\ \bar{\nabla}_i a_{\alpha|j}^k &= \left(\nabla_i a_{\alpha|j}^k - \Gamma_{\alpha|i}^{\perp\beta} a_{\beta|j}^k\right), \quad \left(\bar{\nabla}_{\frac{\partial}{\partial u^i}} A\right)_{n_{\alpha|}}\left(\frac{\partial}{\partial u^j}\right) = \bar{\nabla}_i a_{\alpha|j}^k \frac{\partial}{\partial u^k}, \end{aligned}$$

where $\|g^{kt}\|$ and $\|\delta^{\alpha\beta}\|$ are inverse matrixes to $\|g_{kt}\|$ and $\|\delta_{\alpha\beta}\|$, respectively.

We assume that a Riemannian manifold M^{n+p} is almost Hermitian manifold with almost complex structure J (see [3], chapter 6, section 6.1). Then M^{n+p} has even dimension: $n+p = 2(m+l)$, where a number $m+l$ is called *complex dimension* of M^{n+p} ; the Riemannian metric $\tilde{g}$ is almost Hermitian, i.e. for any vector fields $\tilde{X}, \tilde{Y}$, tangent to M^{n+p} , the following condition holds:

$$\tilde{g}(J\tilde{X}, J\tilde{Y}) = \tilde{g}(\tilde{X}, \tilde{Y}). \quad (1.10)$$

Almost Hermitian manifold M^{n+p} is called *Kaehler manifold* ([3]) if almost complex structure J is parallel, i.e. for any vector fields $\tilde{X}, \tilde{Y}$, tangent to M^{n+p} , the following condition holds:

$$\widetilde{\nabla}_{\tilde{X}} J\tilde{Y} = J\widetilde{\nabla}_{\tilde{X}} \tilde{Y}. \quad (1.11)$$

A submanifold F^n of a Kaehler manifold M^{n+p} is called *Kaehler submanifold* if for any vector field $X \in TF^n$, vector field $JX \in TF^n$. F^n is Kaehler manifold relative to induced almost complex structure J and induced almost Hermitian metric g (see [3], chapter 6, par. 6.7). Kaehler submanifold F^n in Kaehler manifold M^{n+p} , has even dimension $n = 2m$ and codimension $p = 2l$. Number m is called *complex dimension*, and number l is called *complex codimension* of Kaehler submanifold F^n .

We denote by $M^{2m+2l}(c)$, a Kaehler manifold of complex dimension $m+l$ of constant holomorphic sectional curvature c . The tensor of Riemannian curvature $\tilde{R}$ of space $M^{2m+2l}(c)$ complies with the formula [1] :

$$\tilde{R}(\tilde{X}, \tilde{Y})\tilde{Z} = \frac{c}{4} \left(\tilde{g}(\tilde{Y}, \tilde{Z})\tilde{X} - \tilde{g}(\tilde{X}, \tilde{Z})\tilde{Y} + \tilde{g}(J\tilde{Y}, \tilde{Z})J\tilde{X} - \tilde{g}(J\tilde{X}, \tilde{Z})J\tilde{Y} + 2\tilde{g}(\tilde{X}, J\tilde{Y})J\tilde{Z} \right), \quad (1.12)$$

for any vector fields $\tilde{X}, \tilde{Y}, \tilde{Z}$, tangent to $M^{2m+2l}(c)$.

2 The properties of covariant derivative $\bar{\nabla}$.

Lemma 1 . Let F^n be a submanifold in a Riemannian manifold M^{n+p} . Then the following equality holds:

$$\tilde{g}((\bar{\nabla}_Z A)_\xi X, Y) = \tilde{g}((\bar{\nabla}_Z b)(X, Y), \xi) \quad \forall X, Y, Z \in TF^n, \quad \forall \xi \in T^\perp F^n. \quad (2.1)$$

Proof. We will find the expressions of the left and the right parts of the equality (2.1), in local coordinates. We assume

$$Z = Z^i \frac{\partial}{\partial u^i}, \quad X = X^j \frac{\partial}{\partial u^j}, \quad Y = Y^k \frac{\partial}{\partial u^k}, \quad \xi = \xi^\alpha n_{\alpha|}. \quad (2.2)$$

We have:

$$\begin{aligned} \tilde{g}((\bar{\nabla}_Z A)_\xi X, Y) &= Z^i X^j Y^k \xi^\alpha g_{sk} \bar{\nabla}_i a_{\alpha|j}^s = Z^i X^j Y^k (\xi^\alpha g_{sk} \nabla_i a_{\alpha|j}^s - \xi^\alpha g_{sk} \Gamma_{\alpha|i}^{\perp\beta} a_{\beta|j}^s) = \\ &= Z^i X^j Y^k (\xi^\alpha \nabla_i (g_{sk} a_{\alpha|j}^s) - \xi^\alpha \Gamma_{\alpha|i}^{\perp\beta} g_{sk} a_{\beta|j}^s) = Z^i X^j Y^k (\xi^\alpha \nabla_i b_{\alpha|jk} - \xi^\alpha \Gamma_{\alpha|i}^{\perp\beta} b_{\beta|jk}) = \\ &= Z^i X^j Y^k (\xi^\alpha \nabla_i (\delta_{\alpha\beta} b_{jk}^\beta) - \xi^\alpha \Gamma_{\alpha|i}^{\perp\beta} \delta_{\beta\gamma} b_{jk}^\gamma) = Z^i X^j Y^k (\xi^\alpha \delta_{\alpha\beta} \nabla_i b_{jk}^\beta - \xi^\alpha \Gamma_{\alpha|i}^{\perp\beta} \delta_{\beta\gamma} b_{jk}^\gamma) = \\ &= Z^i X^j Y^k (\xi^\alpha \delta_{\alpha\beta} \nabla_i b_{jk}^\beta - \xi^\alpha \Gamma_{\gamma\alpha|i}^\perp b_{jk}^\gamma) = \\ &= Z^i X^j Y^k (\xi^\alpha \delta_{\alpha\beta} (\bar{\nabla}_i b_{jk}^\beta - \Gamma_{\gamma|i}^{\perp\beta} b_{jk}^\gamma) - \xi^\alpha \Gamma_{\gamma\alpha|i}^\perp b_{jk}^\gamma) = \\ &= Z^i X^j Y^k (\xi^\alpha \delta_{\alpha\beta} \bar{\nabla}_i b_{jk}^\beta - \xi^\alpha \delta_{\alpha\beta} \Gamma_{\gamma|i}^{\perp\beta} b_{jk}^\gamma - \xi^\alpha \Gamma_{\gamma\alpha|i}^\perp b_{jk}^\gamma) = \\ &= Z^i X^j Y^k (\xi^\alpha \delta_{\alpha\beta} \bar{\nabla}_i b_{jk}^\beta - \xi^\alpha \Gamma_{\alpha\gamma|i}^\perp b_{jk}^\gamma - \xi^\alpha \Gamma_{\gamma\alpha|i}^\perp b_{jk}^\gamma) = Z^i X^j Y^k \delta_{\alpha\beta} \xi^\alpha \bar{\nabla}_i b_{jk}^\beta = \\ &= \tilde{g}((\bar{\nabla}_Z b)(X, Y), \xi). \end{aligned}$$

Lemma is proved.

Lemma 2 . Let F^{2m} be a Kaehler submanifold in a Kaehler manifold M^{2m+2l} . Then for any $X \in TF^{2m}$ and for any $\xi \in T^\perp F^{2m}$ the following equality holds:

$$(\bar{\nabla}_X A)_{J\xi} = J(\bar{\nabla}_X A)_\xi \quad (2.3)$$

Proof. From (1.1), because of (1.11), we obtain the following equalities (see , for example, [3] , chapter 6, section 6.1, lemma 6. 26):

$$\nabla_X JY = J\nabla_X Y, \quad Jb(X, Y) = b(X, JY), \quad \forall X, Y \in TF^{2m}. \quad (2.4)$$

From (1.2) we have:

$$\widetilde{\nabla}_X J\xi = -A_{J\xi} X + D_X J\xi, \quad J\widetilde{\nabla}_X \xi = J(-A_\xi X + D_X \xi).$$

Hence, because of (1.11), we obtain:

$$-A_{J\xi}X + D_X J\xi = J(-A_\xi X + D_X \xi).$$

Therefore,

$$-A_{J\xi}X + JA_\xi X = JD_X \xi - D_X J\xi.$$

Since F^{2m} is a Kaehler submanifold, then, from here, we have

$$A_{J\xi}X = JA_\xi X, \quad D_X(J\xi) = JD_X \xi, \quad \forall X, Y \in TF^{2m}. \quad (2.5)$$

From (1.7) we have

$$(\bar{\nabla}_X A)_{J\xi}Y = \nabla_X(A_{J\xi}Y) - A_{J\xi}(\nabla_X Y) - A_{D_X(J\xi)}Y.$$

Hence, using (2.4) and (2.5), we have:

$$\begin{aligned} (\bar{\nabla}_X A)_{J\xi}Y &= \nabla_X J(A_\xi Y) - JA_\xi(\nabla_X Y) - A_{J(D_X \xi)}Y = \\ &= J\nabla_X(A_\xi Y) - JA_\xi(\nabla_X Y) - JA_{D_X \xi}Y = J(\bar{\nabla}_X A)_\xi Y. \end{aligned}$$

Lemma is proved.

Lemma 3 . *Let F^{2m} be a Kaehler submanifold in a Kaehler manifold $M^{2m+2l}(c)$ of constant holomorphic sectional curvature c . Then for any $X, Y, Z \in TF^{2m}$ and for any $\xi \in T^\perp F^{2m}$ the following equalities hold:*

$$(\bar{\nabla}_{JZ}b)(X, Y) = J\left((\bar{\nabla}_Z b)(X, Y)\right), \quad (2.6)$$

$$(\bar{\nabla}_{JZ}A)_\xi = -J(\bar{\nabla}_Z A)_\xi, \quad (2.7)$$

$$JA_\xi = -A_\xi J, \quad (2.8)$$

$$J(\bar{\nabla}_Z A)_\xi = -(\bar{\nabla}_Z A)_\xi J. \quad (2.9)$$

Proof. 1. Because of (1.12), the equation (1.4) takes the following form:

$$(\bar{\nabla}_X b)(Y, Z) = (\bar{\nabla}_Y b)(X, Z), \quad \forall X, Y, Z \in TF^{2m}. \quad (2.10)$$

Using (2.10), from (1.7) we obtain:

$$(\bar{\nabla}_{JZ}b)(X, Y) = (\bar{\nabla}_X b)(JZ, Y) = D_X(b(JZ, Y)) - b(\nabla_X(JZ), Y) - b(JZ, \nabla_X Y).$$

Hence, using (2.4) and (2.5), we have:

$$\begin{aligned} (\bar{\nabla}_{JZ}b)(X, Y) &= D_X(J(b(Z, Y))) - b(J\nabla_X Z, Y) - b(JZ, \nabla_X Y) = \\ &= J(D_X(b(Z, Y))) - J(b(\nabla_X Z, Y)) - J(b(Z, \nabla_X Y)) = \end{aligned}$$

$$= J(D_X(b(Z, Y)) - b(\nabla_X Z, Y) - b(Z, \nabla_X Y)) = J((\bar{\nabla}_Z b)(X, Y)).$$

The equality (2.6) is proved.

2. Using (2.6), from (2.1) we obtain:

$$\tilde{g}((\bar{\nabla}_{JZ} A)_\xi X, Y) = \tilde{g}((\bar{\nabla}_{JZ} b)(X, Y), \xi) = \tilde{g}(J((\bar{\nabla}_Z b)(X, Y)), \xi).$$

Hence, because of (1.10) and equality $J^2 = -I$, we have:

$$\tilde{g}((\bar{\nabla}_{JZ} A)_\xi X, Y) = -\tilde{g}((\bar{\nabla}_Z b)(X, Y), J\xi) = -\tilde{g}((\bar{\nabla}_Z A)_{J\xi} X, Y) = -\tilde{g}(J((\bar{\nabla}_Z A)_\xi X), Y).$$

From here we get (2.7).

3. From (1.6), using (2.4), we obtain:

$$\tilde{g}(JA_\xi X, Y) = -\tilde{g}(A_\xi X, JY) = -\tilde{g}(b(X, JY), \xi) = -\tilde{g}(b(JX, Y), \xi) = -\tilde{g}(A_\xi JX, Y).$$

Thus,

$$\tilde{g}(JA_\xi X, Y) = -\tilde{g}(A_\xi JX, Y) \quad \forall X, Y \in TF^{2m}, \quad \forall \xi \in T^\perp F^{2m}.$$

The derived equality is equivalent to (2.8).

4. From (1.8), using (2.4) and (2.8), for any $X, Y \in TF^{2m}$ and for any $\xi \in T^\perp F^{2m}$, we have:

$$\begin{aligned} J((\bar{\nabla}_X A)_\xi Y) &= J(\nabla_X(A_\xi Y) - A_\xi(\nabla_X Y) - A_{D_X \xi} Y) = \\ &= \nabla_X J(A_\xi Y) + A_\xi J(\nabla_X Y) + A_{D_X \xi}(JY) = \\ &= -\nabla_X(A_\xi JY) + A_\xi(\nabla_X JY) + A_{D_X \xi}(JY) = -(\bar{\nabla}_X A)_\xi(JY). \end{aligned}$$

Thus,

$$J((\bar{\nabla}_X A)_\xi Y) = -(\bar{\nabla}_X A)_\xi(JY), \quad \forall X, Y \in TF^{2m}, \quad \forall \xi \in T^\perp F^{2m}.$$

The obtained equality is equivalent to (2.9). Lemma is proved.

Lemma 4 . *Let F^{2m} be a Kaehler submanifold in a Kaehler manifold M^{2m+2l} . Then the following equality holds*

$$\bar{\nabla}_Z(\tilde{g}(X, JY)J\xi) = 0 \quad \forall X, Y, Z \in TF^{2m}, \quad \forall \xi \in T^\perp F^{2m}. \quad (2.11)$$

Proof. By the definition of covariant derivative $\bar{\nabla}$, we have:

$$\begin{aligned} \bar{\nabla}_Z(\tilde{g}(X, JY)J\xi) &= \\ &= D_Z(\tilde{g}(X, JY)J\xi) - \tilde{g}(\nabla_Z X, JY)J\xi - \tilde{g}(X, \nabla_Z(JY))J\xi - \tilde{g}(X, JY)D_Z(J\xi). \end{aligned}$$

We transform the right part of the last equality, writing it in local coordinates and using the designations (2.2):

$$\left(\frac{\partial (g_{kl} X^k (JY)^l (J\xi)^\tau)}{\partial u^i} + \Gamma_{\sigma|i}^\perp g_{kl} X^k (JY)^l (J\xi)^\sigma \right) Z^i n_\tau -$$

$$\begin{aligned}
& -g_{kl} \left(\frac{\partial X^k}{\partial u^i} + \Gamma_{im}^k X^m \right) (JY)^l (J\xi)^\tau Z^i n_\tau - g_{lk} \left(\frac{\partial (JY)^l}{\partial u^i} + \Gamma_{im}^l (JY)^m \right) X^k (J\xi)^\tau Z^i n_\tau - \\
& -g_{kl} X^k (JY)^l \left(\frac{\partial (J\xi)^\tau}{\partial u^i} + \Gamma_{\sigma|i}^\tau (J\xi)^\sigma \right) Z^i n_\tau = \\
& = \left(\frac{\partial g_{kl}}{\partial u^i} X^k (JY)^l (J\xi)^\tau - g_{kl} \Gamma_{im}^k X^m (JY)^l (J\xi)^\tau - g_{lk} \Gamma_{im}^l X^k (JY)^m (J\xi)^\tau \right) Z^i n_\tau = \\
& = \left(\frac{\partial g_{kl}}{\partial u^i} - g_{ml} \Gamma_{ik}^m - g_{mk} \Gamma_{il}^m \right) X^k (JY)^l (J\xi)^\tau Z^i n_\tau = 0.
\end{aligned}$$

Lemma is proved.

Lemma 5 . *Let F^{2m} be a Kaehler submanifold in a Kaehler manifold $M^{2m+2l}(c)$ of constant holomorphic sectional curvature c . Then the following equality holds*

$$\begin{aligned}
R^\perp(X, Y)\xi &= \frac{c}{2} \tilde{g}(X, JY) J\xi + b(X, A_\xi Y) - b(Y, A_\xi X), \\
\forall X, Y &\in TF^{2m}, \quad \forall \xi \in T^\perp F^{2m}.
\end{aligned} \tag{2.12}$$

Proof. Because of (1.12), we have:

$$\tilde{R}(X, Y, \xi, \eta) = \tilde{g}(\tilde{R}(X, Y)\xi, \eta) = \frac{c}{2} \tilde{g}(X, JY) \tilde{g}(J\xi, \eta).$$

Then the equation (1.5) takes the form:

$$R^\perp(X, Y, \xi, \eta) = \frac{c}{2} \tilde{g}(X, JY) \tilde{g}(J\xi, \eta) + \tilde{g}([A_\xi, A_\eta]X, Y).$$

We transform the second term in the right part of the obtained equality, using self-adjointness of operator A_ξ :

$$\begin{aligned}
\tilde{g}([A_\xi, A_\eta]X, Y) &= \tilde{g}((A_\xi A_\eta - A_\eta A_\xi)X, Y) = \tilde{g}(A_\xi(A_\eta X), Y) - \tilde{g}(A_\eta(A_\xi X), Y) = \\
&= \tilde{g}(A_\eta X, A_\xi Y) - \tilde{g}(A_\xi X, A_\eta Y) = \tilde{g}(b(A_\xi Y, X), \eta) - \tilde{g}(b(A_\xi X, Y), \eta).
\end{aligned}$$

Then for any $\eta \in T^\perp F^{2m}$ we have:

$$\begin{aligned}
R^\perp(X, Y, \xi, \eta) &\equiv \tilde{g}(R^\perp(X, Y)\xi, \eta) = \\
&= \tilde{g}\left(\frac{c}{2} \tilde{g}(X, JY) J\xi, \eta\right) + \tilde{g}(b(A_\xi Y, X), \eta) - \tilde{g}(b(A_\xi X, Y), \eta).
\end{aligned}$$

From here we obtain the equality (2.12). Lemma is proved.

Lemma 6 . Let F^{2m} be a Kaehler submanifold in a Kaehler manifold $M^{2m+2l}(c)$ of constant holomorphic sectional curvature c . Then the following equality holds

$$\begin{aligned} & (\bar{\nabla}_Z R^\perp)(X, Y)\xi = \\ & = (\bar{\nabla}_Z b)(X, A_\xi Y) + b(X, (\bar{\nabla}_Z A)_\xi Y) - (\bar{\nabla}_Z b)(Y, A_\xi X) - b(Y, (\bar{\nabla}_Z A)_\xi X), \\ & \quad \forall X, Y, Z \in TF^{2m}, \quad \forall \xi \in T^\perp F^{2m}. \end{aligned} \quad (2.13)$$

Proof. From formula (1.9), using (2.12), we obtain:

$$\begin{aligned} & (\bar{\nabla}_Z R^\perp)(X, Y)\xi = D_Z \left(\frac{c}{2} \tilde{g}(X, JY) J\xi + b(X, A_\xi Y) - b(Y, A_\xi X) \right) - \\ & \quad - \left(\frac{c}{2} \tilde{g}(\nabla_Z X, JY) J\xi + b(\nabla_Z X, A_\xi Y) - b(Y, A_\xi(\nabla_Z X)) \right) - \\ & \quad - \left(\frac{c}{2} \tilde{g}(X, J(\nabla_Z Y)) J\xi + b(X, A_\xi(\nabla_Z Y)) - b(\nabla_Z Y, A_\xi X) \right) - \\ & \quad - \left(\frac{c}{2} \tilde{g}(X, JY) J(D_Z \xi) + b(X, A_{D_Z \xi} Y) - b(Y, A_{D_Z \xi} X) \right) = \\ & = \frac{c}{2} \left(D_Z(\tilde{g}(X, JY) J\xi) - \tilde{g}(\nabla_Z X, JY) J\xi - \tilde{g}(X, J(\nabla_Z Y)) J\xi - \tilde{g}(X, JY) J(D_Z \xi) \right) + \\ & \quad + D_Z(b(X, A_\xi Y)) - D_Z(b(Y, A_\xi X)) - b(\nabla_Z X, A_\xi Y) + b(Y, A_\xi(\nabla_Z X)) - \\ & \quad - b(X, A_\xi(\nabla_Z Y)) + b(\nabla_Z Y, A_\xi X) - b(X, A_{D_Z \xi} Y) + b(Y, A_{D_Z \xi} X). \end{aligned}$$

Hence, using (2.3) and (2.4), we have

$$\begin{aligned} & (\bar{\nabla}_Z R^\perp)(X, Y)\xi = \\ & = \frac{c}{2} \bar{\nabla}_Z(\tilde{g}(X, JY) J\xi) + D_Z(b(X, A_\xi Y)) - D_Z(b(Y, A_\xi X)) - b(\nabla_Z X, A_\xi Y) + \\ & \quad + b(Y, A_\xi(\nabla_Z X)) - b(X, A_\xi(\nabla_Z Y)) + b(\nabla_Z Y, A_\xi X) - b(X, A_{D_Z \xi} Y) + b(Y, A_{D_Z \xi} X). \end{aligned}$$

Therefore, because of (2.11), we obtain:

$$\begin{aligned} & (\bar{\nabla}_Z R^\perp)(X, Y)\xi = \left(D_Z(b(X, A_\xi Y)) - b(\nabla_Z X, A_\xi Y) - b(X, A_\xi(\nabla_Z Y)) - b(X, A_{D_Z \xi} Y) \right) - \\ & \quad - \left(D_Z(b(Y, A_\xi X)) - b(\nabla_Z Y, A_\xi X) - b(Y, A_\xi(\nabla_Z X)) - b(Y, A_{D_Z \xi} X) \right). \end{aligned}$$

Hence, using (1.7), we obtain:

$$(\bar{\nabla}_Z R^\perp)(X, Y)\xi = \left((\bar{\nabla}_Z b)(X, A_\xi Y) + b(X, \nabla_Z(A_\xi Y)) - b(X, A_\xi(\nabla_Z Y)) - b(X, A_{D_Z \xi} Y) \right) -$$

$$-\left((\bar{\nabla}_Z b)(Y, A_\xi X) + b(Y, \nabla_Z(A_\xi X)) - b(Y, A_\xi(\nabla_Z X)) - b(Y, A_{D_Z \xi} X)\right).$$

Now, using (1.8), we obtain:

$$\begin{aligned} & (\bar{\nabla}_Z R^\perp)(X, Y)\xi = \\ & = \left((\bar{\nabla}_Z b)(X, A_\xi Y) + b(X, (\bar{\nabla}_Z A)_\xi Y)\right) - \left((\bar{\nabla}_Z b)(Y, A_\xi X) + b(Y, (\bar{\nabla}_Z A)_\xi X)\right). \end{aligned}$$

Lemma is proved.

Lemma 7 . Let F^{2m} be a Kaehler submanifold in a Kaehler manifold $M^{2m+2l}(c)$ of constant holomorphic sectional curvature c . Then the following equality holds

$$\begin{aligned} (\bar{\nabla}_Z R^\perp)(X, Y, \xi, \eta) &= \tilde{g}([\bar{\nabla}_Z A]_\xi, A_\eta]X, Y) + \tilde{g}([A_\xi, (\bar{\nabla}_Z A)_\eta]X, Y), \\ \forall X, Y, Z \in TF^{2m}, \quad \forall \xi, \eta \in T^\perp F^{2m}. \end{aligned} \quad (2.14)$$

Proof. Because of (2.13), we have:

$$\begin{aligned} (\bar{\nabla}_Z R^\perp)(X, Y, \xi, \eta) &\equiv \tilde{g}\left((\bar{\nabla}_Z R^\perp)(X, Y)\xi, \eta\right) = \tilde{g}\left((\bar{\nabla}_Z b)(X, A_\xi Y), \eta\right) - \\ &- \tilde{g}\left((\bar{\nabla}_Z b)(Y, A_\xi X), \eta\right) + \tilde{g}\left(b(X, (\bar{\nabla}_Z A)_\xi Y), \eta\right) - \tilde{g}\left(b(Y, (\bar{\nabla}_Z A)_\xi X), \eta\right). \end{aligned}$$

In the derived equality, we transform the first and the second terms using (2.1), the third and the fourth using (1.6):

$$\begin{aligned} (\bar{\nabla}_Z R^\perp)(X, Y, \xi, \eta) &= \tilde{g}\left((\bar{\nabla}_Z A)_\eta X, A_\xi Y\right) - \tilde{g}\left((\bar{\nabla}_Z A)_\eta Y, A_\xi X\right) + \\ &+ \tilde{g}\left(A_\eta X, (\bar{\nabla}_Z A)_\xi Y\right) - \tilde{g}\left(A_\eta Y, (\bar{\nabla}_Z A)_\xi X\right). \end{aligned}$$

Hence, because of self-adjointness of operators A_ξ and $(\bar{\nabla}_Z A)_\xi$, we obtain:

$$\begin{aligned} (\bar{\nabla}_Z R^\perp)(X, Y, \xi, \eta) &= \tilde{g}\left(A_\xi(\bar{\nabla}_Z A)_\eta X, Y\right) - \tilde{g}\left(Y, (\bar{\nabla}_Z A)_\eta A_\xi X\right) + \\ &+ \tilde{g}\left((\bar{\nabla}_Z A)_\xi A_\eta X, Y\right) - \tilde{g}\left(Y, A_\eta(\bar{\nabla}_Z A)_\xi X\right) = \\ &= \tilde{g}\left([A_\xi, (\bar{\nabla}_Z A)_\eta]X, Y\right) + \tilde{g}\left([\bar{\nabla}_Z A]_\xi, A_\eta]X, Y\right). \end{aligned}$$

Lemma is proved.

Lemma 8 . Let F^{2m} be a Kaehler submanifold in a Kaehler manifold $M^{2m+2l}(c)$ of constant holomorphic sectional curvature c . Then the following equality holds

$$\begin{aligned} (\bar{\nabla}_{JZ} R^\perp)(X, Y, \xi, \eta) &= (\bar{\nabla}_Z R^\perp)(X, Y, J\xi, \eta) - 2\tilde{g}([\bar{\nabla}_Z A]_{J\xi}, A_\eta]X, Y), \\ \forall X, Y, Z \in TF^{2m}, \quad \forall \xi, \eta \in T^\perp F^{2m}. \end{aligned} \quad (2.15)$$

Proof. From (2.14) we obtain:

$$(\bar{\nabla}_{JZ}R^\perp)(X, Y, \xi, \eta) = \tilde{g}([\bar{\nabla}_{JZ}A)_\xi, A_\eta]X, Y) + \tilde{g}([A_\xi, (\bar{\nabla}_{JZ}A)_\eta]X, Y).$$

Hence, using (2.7), we have:

$$(\bar{\nabla}_{JZ}R^\perp)(X, Y, \xi, \eta) = \tilde{g}([-J(\bar{\nabla}_ZA)_\xi, A_\eta]X, Y) + \tilde{g}([A_\xi, -J(\bar{\nabla}_ZA)_\eta]X, Y).$$

In the derived equality, we transform the second term using (2.8) and (2.9):

$$\begin{aligned} [A_\xi, J(\bar{\nabla}_ZA)_\eta] &= A_\xi J(\bar{\nabla}_ZA)_\eta - J(\bar{\nabla}_ZA)_\eta A_\xi = \\ &= -JA_\xi(\bar{\nabla}_ZA)_\eta + (\bar{\nabla}_ZA)_\eta JA_\xi = -[JA_\xi, (\bar{\nabla}_ZA)_\eta]. \end{aligned}$$

Therefore,

$$(\bar{\nabla}_{JZ}R^\perp)(X, Y, \xi, \eta) = -\tilde{g}([J(\bar{\nabla}_ZA)_\xi, A_\eta]X, Y) + \tilde{g}([JA_\xi, (\bar{\nabla}_ZA)_\eta]X, Y).$$

Hence, because of (2.3) and (2.5), and using (2.14), we obtain:

$$\begin{aligned} (\bar{\nabla}_{JZ}R^\perp)(X, Y, \xi, \eta) &= -\tilde{g}([\bar{\nabla}_ZA)_{J\xi}, A_\eta]X, Y) + \tilde{g}([A_{J\xi}, (\bar{\nabla}_ZA)_\eta]X, Y) = \\ &= (\bar{\nabla}_ZR^\perp)(X, Y, J\xi, \eta) - 2\tilde{g}([\bar{\nabla}_ZA)_{J\xi}, A_\eta]X, Y). \end{aligned}$$

Lemma is proved.

3 Proofs of theorems 1, 2.

Proof theorem 1.

Let for some 1-form μ on F^{2m} , the following condition holds

$$(\bar{\nabla}_X b)(Y, Z) = \mu(X)b(Y, Z) \quad \forall X, Y, Z \in TF^{2m}. \quad (3.1)$$

Then for any vector field $\xi \in T^\perp F^{2m}$, we have:

$$\tilde{g}((\bar{\nabla}_X b)(Y, Z), \xi) = \tilde{g}(\mu(X)b(Y, Z), \xi).$$

Hence, using (2.1) and (1.6), we obtain:

$$\tilde{g}((\bar{\nabla}_X A)_\xi Y, Z) = \tilde{g}(\mu(X)A_\xi Y, Z) \quad \forall X, Y, Z \in TF^{2m}, \quad \forall \xi \in T^\perp F^{2m}.$$

Thus, the condition (3.1) is equivalent to the condition

$$(\bar{\nabla}_X A)_\xi = \mu(X)A_\xi, \quad \forall X \in TF^{2m}, \quad \forall \xi \in T^\perp F^{2m}. \quad (3.2)$$

From (3.2) we obtain the equality:

$$(\bar{\nabla}_{JX} A)_\xi = \mu(JX)A_\xi \quad \forall X \in TF^{2m}, \quad \forall \xi \in T^\perp F^{2m}. \quad (3.3)$$

On the other hand, from (3.2), because of (2.7), we have:

$$\left(\bar{\nabla}_{JX}A\right)_{\xi} = -J(\mu(X)A_{\xi}), \quad \forall X \in TF^{2m}, \quad \forall \xi \in T^{\perp}F^{2m}. \quad (3.4)$$

From (3.3) and (3.4), we obtain:

$$\mu(JX)A_{\xi} = -J(\mu(X)A_{\xi}), \quad \forall X \in TF^{2m}, \quad \forall \xi \in T^{\perp}F^{2m}.$$

Hence, for any $Y \in TF^{2m}$, we have:

$$\mu(JX)A_{\xi}Y = -\mu(X)J(A_{\xi}Y), \quad \forall X \in TF^{2m}, \quad \forall \xi \in T^{\perp}F^{2m}. \quad (3.5)$$

Using (3.5), we obtain:

$$\begin{aligned} \mu(JX)\tilde{g}(A_{\xi}Y, A_{\xi}Y) &= -\mu(X)\tilde{g}(J(A_{\xi}Y), A_{\xi}Y) = 0, \\ \forall X, Y \in TF^{2m}, \quad \forall \xi \in T^{\perp}F^{2m}. \end{aligned} \quad (3.6)$$

Since $b \neq 0$ then there exists nondegenerate vector field $\xi \in T^{\perp}F^{2m}$, and from (3.6) we come to the equality:

$$\mu(X) = 0 \quad \forall X \in TF^{2m}.$$

Then 1-form $\mu \equiv 0$ and, therefore,

$$\left(\bar{\nabla}_X A\right)_{\xi} = 0, \quad \forall X \in TF^{2m}, \quad \forall \xi \in T^{\perp}F^{2m}. \quad (3.7)$$

Hence, because of (2.14), we obtain the conclusion of the theorem.

Proof of theorem 2.

Form (1.3) we obtain:

$$\begin{aligned} \nabla_W R(X, Y, Z, V) &= \tilde{g}((\bar{\nabla}_W b)(X, V), b(Y, Z)) + \tilde{g}(b(X, V), (\bar{\nabla}_W b)(Y, Z)) - \\ &- \tilde{g}((\bar{\nabla}_W b)(X, Z), b(Y, V)) - \tilde{g}(b(X, Z), (\bar{\nabla}_W b)(Y, V)) \quad \forall X, Y, Z, V, W \in TF^{2m}. \end{aligned}$$

Therefore, because of (3.7), $\nabla R \equiv 0$. Theorem is proved.

References

- [1] Kobayashi S., Nomizu K. Foundations of differential geometry. Vol. 2. M.: Nauka. 1981.
- [2] Chen B.-Y. Geometry of submanifolds. N.-Y.: M. Dekker. 1973.
- [3] Gray A. Tubes. M.: Mir. 1993.